

Honey as a micro-bacterial agent: identification method of the compounds that inhibit pathogenic bacteria

Arias-De la Cruz, Heydi L.¹; Zaldívar-Cruz, Juan M.^{1*}; Ruiz-May, Eliel², Hernández-Nataren, Edith¹, Sol-Sánchez, Ángel¹, Nila-Méndez, Alejandro G.³

¹ Colegio de Postgraduados - Campus Tabasco, Periférico Carlos A. Molina, Tabasco, México, C. P. 86500.

² Instituto de Ecología A.C., Colonia El Haya, Xalapa, Veracruz, México, C. P. 91073.

³ Tecnológico Nacional de México/ITS Acatlán, Colonia Agrícola Michapa, Acatlán, Veracruz, México, C. P. 96100.

* Correspondence: zaldivar@colpos.mx

ABSTRACT

Objective: To provide an overview of the scientific evidence that supports the use of *Apis mellifera* honey as an antibacterial agent.

Design/Methodology/Approach: An exhaustive review of scientific literature was carried out. The collected information included the different honey types that, according to the reports, have antibacterial properties. In addition, the related compounds, the main chromatographic methods used for their identification, and the main pathogens that have been studied were analyzed.

Results: The antibacterial properties of honey (especially monofloral honeys) have been widely studied worldwide, focusing on their capacity to inhibit pathogenic bacteria. The different methods used to study honey include agar diffusion, disk diffusion, and broth and agar dilution. These properties have been attributed to honey, as a result of its high sugar content, low moisture content, and acidic pH, as well as the diversity of the chemical compounds—mainly hydrogen peroxide, methylglyoxal (MGO), phenolic acids, flavonoids, peptides, glycopeptides, and different proteins—that were identified by a chromatographic analysis.

Study Limitations/Implications: Currently, the honey of bees (*Apis mellifera*) has great potential as an alternative to combat the antibiotic resistance of certain pathogens.

Findings/Conclusions: Honey can inhibit both gram-positive and -negative bacteria, such as *Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, and *Enterobacter*. These characteristics are the result of the diverse chemical compounds of honey. In addition, these compounds widely change depending on the vegetation that surrounds the hives; therefore, honey from different geographical origins has unique characteristics, in terms of its composition and antibacterial activity.

Keywords: *Apis mellifera*, honey, antibacterial property.

INTRODUCTION

Honey is a sweet substance produced by bees, from the flower nectar or exudations of live parts of plants, that is transformed and stored in their honeycomb (Martínez-González and Pérez-López, 2013). This natural product consists of approximately 38% fructose, 31% glucose, 10% of different types of sugar, 18% water, and 3% of other compounds—such as

Citation: Arias-De la Cruz, H. L., Zaldivar-Cruz, J. M., Ruiz-May, E., Hernández-Nataren, E., Sol-Sánchez, Á., & Nila-Méndez, A. G. (2024) Honey as a micro-bacterial agent: identification method of the compounds that inhibit pathogenic bacteria. *Agro Productividad*. <https://doi.org/10.32854/agrop.v17i8.2631>

Academic Editor: Jorge Cadena Iñiguez

Guest Editor: Juan Francisco Aguirre Medina

Received: July 10, 2023.

Accepted: June 28, 2024.

Published on-line: September 02, 2024.

Agro Productividad, 17(8). August 2024.
pp: 35-52.

This work is licensed under a
Creative Commons Attribution-Non-
Commercial 4.0 International license.



proteins, lipids, amino acids, phenolic compounds, minerals, vitamins, and carotenoids (Hegazi *et al.*, 2014; Bueno-Costa *et al.*, 2016; Visweswara *et al.*, 2016; Deng *et al.*, 2018; Cheung *et al.*, 2019; Leyva-Jiménez *et al.*, 2019).

Honey is very important worldwide and it has been used to treat skin wounds, burns, sores, eye infections, sore throat, and other issues. The properties of honey are the consequence of its high thickness that protects against infections, its enzymatic production of hydrogen peroxide (H_2O_2), its high osmolarity, and its low pH (3.2-4.5). As a whole, these characteristics inhibit the growth of pathogenic bacteria (Mandal and Mandal, 2011). In addition, honey is consumed as a result of its high nutritional value and its antioxidant, bacteriostatic, and anti-inflammatory properties (Alvarez-Suarez *et al.*, 2014); these biological compounds are passed from the nectar to the honey and can be found in high levels (Güneş *et al.*, 2017).

The inappropriate use of antibiotics to treat bacterial infections has produced different degrees of bacterial resistance and, consequently, the use of these products has been limited. *Escherichia coli*, *Staphylococcus aureus*, and *Salmonella enterica* are some of the hospital pathogens that cause difficult-to-treat infections; they have developed an increasing resistance to antibiotics, causing a worldwide concern (Da Silva and Mendonça, 2012; Fyfe *et al.*, 2017).

Consequently, finding new alternatives to improve human health is fundamental. Some of these alternatives include the combination of antibacterial formulations aimed to produce more efficient antibiotics or the search for natural alternatives, such as apitherapy. This last alternative offers honey-based treatments and other honeycomb products that can be used to combat several bacterial infections (Estrada *et al.*, 2005; Mandal and Mandal, 2011; Boussaid *et al.*, 2018). Jenkins and Cooper (2012) have reported that the oxacillin and Manuka honey synergy can be used to treat *S. aureus*, which is resistant to oxacillin. In addition, the Manuka honey and rifampicin (antibiotic) combination can be used to inhibit *S. aureus*, which is resistant to methicillin (MRSA). The antibiotic resistance reversion is a consequence of the reduction of the *mecR1* (MRSA resistance gene) caused by honey (Müller *et al.*, 2013).

Consequently, the objective of this review was to gather scientific evidence about the properties of honey produced by bees (*Apis mellifera*) as an antibacterial agent and to identify the main methods used to establish the secondary metabolites involved in the antibacterial potential.

MATERIALS AND METHODS

A bibliographic review was carried out using the PubMed, Google Scholar, Scopus, and Science Direct databases, in order to gather information about the main methodologies used to evaluate the antibacterial properties of honeys from different geographical origins. In addition, the main secondary metabolites related with the antibacterial activity of honey and the chromatographic methods used to identify them, during the last twenty-five years, were integrated into the review. The search criteria were limited to English, Portuguese, and Spanish and were based on the scientific publications about antibacterial activity of *Apis mellifera* honey, antibacterial metabolites, and antibacterial properties identification

methods. The Mendeley Reference Manager was used to choose and download only those open access publications that included key information.

RESULTS AND DISCUSSION

The initial search identified 250 publications in the different abovementioned databases. Once all the publications were analyzed, 140 were excluded, because their content was not related to the objective of this study. Most of the chosen publications were experimental qualitative studies, which identified both international and domestic honey compounds and the equipment used for this purpose.

Methods and techniques used in the identification of antibacterial compounds

As a result of its many compounds, honey is a complex matrix; however, these compounds can be analyzed using chromatographic techniques. Pyrzynska and Biesaga (2009) tested methods and techniques to obtain higher extraction yields for the different chemical structures and functional groups (aldehydes, ketones, carboxylic acid, esters, alcohols, and flavonoids).

The chromatographic analysis requires a representative sample, which must be subjected to a pre-treatment (extraction) to remove any compound that could interfere in the analysis, such as sugars and polar substances. Highly polar compounds (flavonoids and phenolic acid) should be subjected to a derivatization (chemical changes in the analyte or sample) (Ciulu *et al.*, 2016).

Analytical tools, such as the UV-Vis spectrophotometer, the spectrofluorometer, and ready-to-use H₂O₂ assay kits has been used to measure the H₂O₂ level of honey (Sowa *et al.*, 2017).

Chen *et al.* (2012) evaluated the H₂O₂ content of processed (heat treatment) and unprocessed Australian honeys, measuring the absorbency at 560 nm, and found a 0-1,017 μM concentration. The correlation of the antibacterial activity was higher in the unprocessed samples, because the compound is sensitive to the heat treatment. Likewise, Sowa *et al.* (2017) determined the H₂O₂ content of monofloral clover honeys (Poland), using an absorbency of 540 nm. These samples recorded activity against gram-positive bacteria, with 12.5-25% concentrations.

Poli *et al.* (2018) evaluated the H₂O₂ concentrations of five honey samples, with and without catalase (an enzyme that catalyzes the decomposition of H₂O₂ into oxygen and water), and they recorded that honey without catalase obtained 7-8% minimum inhibitory concentration (MIC) values, proving its high antibacterial capacity. Meanwhile, when these authors added catalase to the samples, the values increased by 25%. These results were corroborated by Sindi *et al.* (2019), who recorded a 29.4% increase with honey treated with catalase.

The H₂O₂ assay kits are very reliable, given their sensibility in presence of peroxidase. The reagent of the kit reacts with H₂O₂, creating resorufin (a fluorescent red oxidation product). This reaction has been used to detect it in <100 μL volumes. Compared with other, cheaper, and easier-to-use colorimeter methods, the main disadvantage of this product is its high cost (Chen *et al.*, 2012; Sowa *et al.*, 2017).

A high-performance liquid chromatography (HPLC) was used to analyze methylglyoxal (MGO) as the quinoxaline content that remained after the o-phenylenediamine derivatization. Atrott and Henle (2009) recorded values that fluctuated between 189 and 835 mg kg⁻¹, showing a good methylglyoxal-antibacterial activity correlation (12.4 and 30.9%, respectively). For their part, Sultanbawa *et al.* (2015) used an ortho-phenylenediamine derivatization and recorded a lineal correlation between the MGO (279-1,755 mg kg⁻¹) content and the *E. coli* and *S. aureus* bacterial inhibition (8-55 mg kg⁻¹ to inactivate the bacteria).

Cokcetin *et al.* (2016) used an O-(2,3,4,5,6-Pentafluorobenzyl) hydroxylamine derivatization and obtained 1,100 mg kg⁻¹, recording a strong correlation between the concentration of the bacterial activity without peroxide and MGO. These results were corroborated by Rückriemen *et al.* (2017).

Other studies have used different methods to identify the phenolic compounds, including a HPLC with a PhotoDiode Array Detector (PDA) and a UV-Vis spectrophotometer. Additionally, they compare the retention time of the analytes with reference standards. Table 1 shows the analytical methods used to identify honey phenolic compounds in different countries. Escriche *et al.* (2014) used a HPLC-PDA to determine the flavonoids and phenolic acids of citrus, rosemary, and honeydew honeys and they identified the following indicators: hesperetin (citrus honey), pinocembrin (rosemary honey), and myricetin and p-coumaric acid (honeydew honey). Güneş *et al.* (2017) found caffeic, protocatechuic, and p-hydroxybenzoic acids in honeys from Turkey; while the most abundant compounds of these honeys were pinocembrin, chrysin, and galangina.

Apigenin, 4-hydroxybenzoic acid, isorhamnetin, luteolin, and pinocembrin were found in all honey samples from different regions of Algeria (Ouchemoukh *et al.*, 2017). Meanwhile, gallic acid (phenolic acid) and chrysin (flavonoids) were the major compounds found in commercial honeys from different countries (Cheung *et al.*, 2019). For their part, Elrasheid *et al.* (2017) detected vanillic, chlorogenic, syringic acid, and catechin in honey samples from Sudan. These bioactive compounds produce antibacterial, antioxidant, and anti-inflammatory effects (Leyva-Jiménez *et al.*, 2019; Goslinski *et al.*, 2020; Velásquez *et al.*, 2020).

Antibacterial properties of honey

The antibacterial properties of honey are the result of its high sugar content and its capacity to generate H₂O₂, produced by the glucose oxidase enzyme synthesized by bees. This compound destroys the essential components of the cells; however, the inhibitory capacity decreases as temperature rises. This phenomenon shows that the antibacterial potential of honey depends on its composition, the nectar source from which bees feed, the harvesting conditions, the pasteurization process, storage time, and temperature of the storing facilities (Fernandes *et al.*, 2020; Goslinski *et al.*, 2020; Velásquez *et al.*, 2020).

The efficiency of H₂O₂ as an antibacterial agent has been proved; nevertheless, in some honeys, it is related to low pH (3.2-4.5), cytokine release, and several molecules with immunomodulatory and anti-inflammatory properties (Almasaudi *et al.*, 2017; Leyva-Jiménez *et al.*, 2019; Fernandes *et al.*, 2020). For example, Melaleuca honey has an acid

Table 1. Main phenolic compounds of honeys from different geographical origins, detected by chromatography.

Country	Secondary metabolites	Honey	Method chromatographic	Reference
New Zealand and y Australia	Myricetin, crysin and chlorogenic, gallic, caffeic, ferulic, <i>p</i> -coumaric, rosmarinic, ellagic, 3,4-dihydroxybenzoic and abscisic acids.	Manuka <i>Leptospermum scoparium</i>	HPLC-DAD, HPTLC	Stanek y Jasicka-Misiak, 2018
Malaysia	Caffeic, chlorogenic, ferulic, <i>p</i> -coumaric, sinapic, vanillic and syringic acids.	<i>Melaleuca alternifolia</i>	RP-UHPLC-ESI-MS	Goslinski <i>et al.</i> , 2021
Poland	Chlorogenic and ferulic acid.	<i>Fagopyrum esculentum</i> , <i>Calluna vulgaris</i> and <i>Tilia</i>	RP-UHPLC-ESI-MS	Goslinski <i>et al.</i> , 2021
Poland	Myricetin, quercetin, naringenin, chrysins and acids: cinnamic, caffeic, abscisic and ferulic.	<i>Robinia pseudoacacia L.</i>	HPTLC	Stanek <i>et al.</i> , 2019
Bangladesh	Catechin, naringin, myricetin, naringenin, hesperetin, kaempferol, apigenin and acids: gallic, chlorogenic, caffeic, coniferous, benzoic and transcinnamic.	<i>Brassica nigra</i> , <i>Nigella sativa</i> , <i>Nelumbo nucifera</i>	HPLC-UV	Moniruzzaman <i>et al.</i> , 2014
Chile	Hymenoptaecin, rutin, naringerin, esculetin, scopoletin, and syringic, <i>p</i> -coumaric, and vanillic acids.	<i>Cryptocarya alba</i> , <i>Quillaja saponaria</i>	HPLC-MS-MS	Montenegro <i>et al.</i> , 2008; Velásquez <i>et al.</i> , 2020
China	Quercetin, apigenin, kaempferol, isorhamnetin, chrysins, galangin and protocatechuic, chlorogenic, caffeic, syringic, <i>p</i> -hydroxybenzoic, <i>p</i> -coumaric, ferulic, isoferulic and benzoic acids.	<i>Fagopyrum esculentum</i>	RP-HPLC-UV	Deng <i>et al.</i> , 2018
Spain	Hesperetin, kaempferol, chrysins, pinocembrin, myricetin, naringenin, quercetin, galangin and caffeic, chlorogenic and <i>p</i> -coumaric acids.	<i>Quercus ilex</i> , <i>Quercus robur</i>	HPLC-PDA,	Escríche <i>et al.</i> , 2014
Portugal	Chrysins, galangin, hesperidin, pinobanksin, luteolin, pinocembrin, kaempferol, apigenin and gallic, caffeic, coumaric and ellagic acids.	<i>Eucalyptus</i> spp., <i>Castanea sativa</i> , <i>Rubus</i> spp., <i>Erica</i> spp.	HPLC-DAD Infinity	Silva <i>et al.</i> , 2020
Saudi Arabia	Apigenin, chrysins, galangin, luteolin, myricetin, naringin and 4-hydroxybenzoic, syringic and gallic acids.	<i>Lavandula dentata</i> , <i>Hypoestes forskaolii</i> , <i>Ziziphus spina-christi</i>	HPLC-PDA	Badjah <i>et al.</i> , 2016
Australia	Apigenin, hesperetin, naringenin neoponcirin, narirutin, biochanin and rosmarinic, chlorogenic and caffeic acids.	<i>Eucalyptus marginata</i> and <i>Corymbia calophylla</i>	HPLC-PDA, HPLC-MS/MS	Tang <i>et al.</i> , 2016; Mani <i>et al.</i> , 2021
Türkiye	Apigenin, pinocembrin, chrysins, galangin, genkwanin and gallic, phydroxybenzoic, caffeic, syringic, salicylic, <i>p</i> -coumaric and transferulic acids.	<i>Castanea sativa</i>	HPLC-DAD	Güneş <i>et al.</i> , 2017
Soudan	Catechin and chlorogenic, syringic, caffeic, <i>p</i> -hydroxybenzoic, vanillic, <i>p</i> -coumaric, ferulic and cinnamic acids.	<i>Acacia nilotica</i> , <i>A. seyal</i> , <i>Z. spina-christi</i> , <i>Amaranthus graecizan</i> , <i>Eucalyptus</i> spp.	FTIR-ATR-PLSR, HPLC-DAD	Elrasheid <i>et al.</i> , 2017
Algeria	Gallic acid, and caffeic acid, chrysins, luteolin, pinocembrin, pinobanksin, myricetin, genistein and daidzein.	<i>Eucalyptus</i> , <i>Ziziphus</i> , <i>Euphorbia</i> , <i>Citrus</i> or <i>Hedysarum</i>	UPLC-PDA-MS/MS	Ouchemoukh <i>et al.</i> , 2017
International commercial honeys	Vanillin, syringaldehyde, protocatechualdehyde and gallic, protocatechual, 2,3,4-trihydroxybenzoic, <i>p</i> -hydroxybenzoic, gentic, chlorogenic, vanillic, caffeic, syringic, <i>p</i> -coumaric, ferulic, synapic, salicylic acids.	<i>Acacia</i> spp., <i>Tilia</i> spp., <i>Leptospermum scoparium</i> , <i>Eriobotrya japonica</i>	RP-HPLC-DAD Infinity	Cheung <i>et al.</i> , 2019

HPLC=High-Performance Liquid Chromatography; PDA=PhotoDiode Array Detector; UV=UV-Vis spectrophotometer; DAD=Diode Array Detector; FTIR=Fourier Transform Infrared Spectroscopy; HPTLC=High-Performance Thin Layer Chromatography; UPLC=Ultra Performance Liquid Chromatography; Ms/Ms (MS-MS)=Tadem Mass Spectrometry; UHPLC=Ultra-High-Performance Liquid Chromatography; RP=Reverse phase; ESI=Electrospray ionization; ATR=Attenuated total reflectance; PLSR=Partial least squares regression.

pH (3.7), perhaps due to the gluconic acid that causes anti-staphylococcal effects (Wen-Jie and Mei-Siew, 2015; Schencke *et al.*, 2016).

The methylglyoxal (MGO) and defensin-1 content of Manuka (*Leptospermum scoparium*) honey results in a high antibacterial activity against *Streptococcus pyogenes*, *Streptococcus mutans*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Enterobacter cloacae*, and *S. aureus* (Irish *et al.*, 2011; Habib *et al.*, 2014; Johnston *et al.*, 2018; Matzen *et al.*, 2018). Consequently, Manuka honey is used as an alternative antibiotic to heal tissue and to reduce microbial infections (Johnston *et al.*, 2018).

Revamil[®], a medical-grade honey, is produced under standardized conditions in greenhouses. Its potent and reproducible antibacterial activity is the result of its osmotic activity, acid pH, and slow and progressive H₂O₂ release. In addition, its MGO content creates favorable and optimal conditions to heal wounds (Kwakman and Zaai, 2012; Pharma GDD, 2022).

Leptospermum honey has high MGO concentrations. Atrott and Henle (2009) and Fernandes *et al.* (2020) have reported up to 800 mg kg⁻¹ MGO content, which is 100 higher than the concentration found in common honey. MGO concentrations increase as the honey matures. This phenomenon is possibly the consequence of the non-enzymatic conversion of trioses that takes place during long-term storage (up to 120 days, at 37 °C). Meanwhile, its antibacterial effect comes from its capacity to inactivate proteins through cross-linking (Adams *et al.*, 2009).

The physiological wound healing processes can be divided into three phases: inflammatory, proliferative, and remodeling. During the inflammatory phase, honey stimulates the monocytes to release inflammatory cytokines. In the proliferative phase, angiogenesis and fibroplasia take place, along with initial re-epithelialization and contraction of the wound. Finally, during the remodeling phase —when collagen is remodeled and realigned along the tension lines—, apoptosis removes the cells that are no longer needed (Schencke *et al.*, 2016).

Polyphenols are major compounds that have been strongly linked to the antibacterial properties of monofloral honeys. In addition, they can be used to determine the floral and geographic origin of honeys (Iqbal *et al.*, 2015; Albaridi, 2019; Mama *et al.*, 2019; Cebrero *et al.*, 2020) and can be found mainly as flavonoids and phenolic acid and its derivatives (Figure 1) (Cauich *et al.*, 2015; Hossen *et al.*, 2017; Leyva-Jiménez *et al.*, 2019).

The identification and quantification of these compounds are fundamental to understand the bioactivity of honey (Habib *et al.*, 2014). Several authors have proved that darker honeys had high levels of phenolic compounds and, at the same time, high pharmacological potential (Brudzynski and Miotti, 2011; Sant'Ana *et al.*, 2014; Sousa *et al.*, 2016). Although individual phenolic compounds have antibacterial activity, their interaction with H₂O₂ should be taken into account, because it can promote this effect (Escuredo *et al.*, 2012; Poli *et al.*, 2018).

The high number of peptides help honey bees to protect themselves from diseases that impact the honeycomb. Honey has four antimicrobial peptide (AMP) families: defensins, apidaecins, abaecins, and hymenotaecin.




Figure 1. Analysis of the enrichment of secondary metabolites identified to date in honey.

Defensins act against gram-negative bacteria, gram-positive bacteria, and fungi. They are responsible for the destruction of the biofilm formed by the bacteria. Reports indicate that defensins amount to $0.04\text{--}5.17 \mu\text{g g}^{-1}$ honey (Kwakman and Zaaij, 2012; Majtan *et al.*, 2012; Dosler and Karaaslan, 2014; Yi *et al.*, 2014; Valachová *et al.*, 2016). Nevertheless, some studies indicate that some gram-negative bacteria are resistant to defensin-1 (Čeřovský and Bém, 2014). Meanwhile, apidaecin is an antibacterial peptide with a high proline content. Once it enters the bacteria through the lipid bilayer, apidaecin stops the protein expression (Li *et al.*, 2012; Larsen *et al.*, 2019).

Abaecin can be found in western honey bees. It inhibits the growth of both gram-negative and gram-positive bacteria through the permeabilization of the bacteria membrane. Its expression and abundance quickly increase in response to a bacterial infection (Randolt *et al.*, 2008; Danihlík *et al.*, 2015; Larsen *et al.*, 2019). Finally, hymenoptaecin has an abundant glycine content that inhibits the growth of gram-positive and gram-negative bacteria (Erban *et al.*, 2019; Larsen *et al.*, 2019).

Antibacterial activity of honey

Different protocols have been used to evaluate the antibacterial activity of honey: agar diffusion test, disk diffusion test, broth and agar dilution method, and gradient methods (E-test and spiral plating technique) (Balouiri *et al.*, 2016; Osés *et al.*, 2016). The most used methods to determine the minimum inhibitory concentrations (MIC) required to

stop or eliminate bacterial growth are the disk diffusion, well diffusion, and broth and agar dilution methods. *S. aureus* is usually the bacterium of choice for tests and *in vitro* antibacterial activity, as a consequence of its resistance to high sugar content and acidity levels. In addition, this bacterium is sensitive to the action of H₂O₂ (Khalil *et al.*, 2014; Osés *et al.*, 2016).

Reproducibility tests are frequently difficult, since the work is carried out with living beings and several methods are used to determine the antibacterial activity of honey (Balouiri *et al.*, 2016; Osés *et al.*, 2016). However, reference material can provide a reliable base for lab work. In addition, reference strains can be used, because mutations can take place or, because depending on the method, the microorganisms can behave differently (Clavijo, 2002; Camaró-Sala *et al.*, 2015; Balouiri *et al.*, 2016; Roshan *et al.*, 2017; Albaridi, 2019; Mama *et al.*, 2019; Rosas *et al.*, 2019).

Chilean honeys, such as the Ulmo (*Eucryphia cordifolia*, Cav.) honey, are drawing attention due to their antibacterial potential. Velásquez *et al.* (2020) evaluated this type of honey using the agar diffusion method to determine their efficiency against *E. coli*, *P. aeruginosa*, and five *S. aureus*-MRSA strains and concluded that their antibacterial activity is mainly the consequence of their phenolic compounds (flavonoids, benzoic acid derivatives, and volatile compounds). Meanwhile, Cebrero *et al.* (2020) used the microdilution technique to evaluate the antibacterial activity of the Peumo (*Cryptocarya alba* (Molina) Loosser) and Quillay (*Quillaja saponaria*, Molina) honeys against *S. aureus* and *P. aeruginosa*, pointing out that the antibacterial activity was the result of their hymenoptaecin content.

Roshan *et al.* (2017) analyzed the antibacterial activity in honey from western Australia. The results were compared with the results obtained by Activon, a medical-grade honey made from Manuka honey. The authors proved that six of the samples inhibited the growth of *S. aureus* and recorded larger inhibition areas than medical-grade honey. Meanwhile, Pasias *et al.* (2018) reported that Greek honeys had high H₂O₂ and hydroxymethylfurfural (HMF) levels. In addition, their acidity and osmolarity resulted in antibacterial activity against *S. aureus*.

For their part, Leyva-Jiménez *et al.* (2019) used the disk diffusion method to extract phenolic compounds from Iranian honeys, testing them against *E. coli*, *P. aeruginosa*, *S. aureus*, and *Enterococcus faecalis*. The authors concluded that individual and collective phenolic compounds do not have the same potential. These results showed a synergy between the different antibacterial compounds.

Hegazi *et al.* (2017) reported that 20 and 30% concentrations of honeys from Saudi Arabia inhibited *S. aureus*, *S. mutans*, *Klebsiella pneumoniae*, *E. coli*, and *P. aeruginosa* strains, as a result of their flower sources, osmotic properties, pH, H₂O₂, flavonoids, and antibacterial volatile substances (*e.g.*, organic acids). Mahendran and Kumarasamy (2015) used the disk diffusion method to evaluate the antibacterial potential of honey collected in Western Ghats, India, during summer and winter, against *S. aureus*, *S. pyogenes*, *E. coli*, *P. aeruginosa*, and *P. mirabilis*. They concluded that the honey samples collected during winter had a higher bactericidal effect than those collected in summer, as a consequence of changes in seasonal flowering.

The eucalyptus honey from the Island of Mauritius is an exceptional case. It showed antibacterial activity against *E. coli*, *P. mirabilis*, *P. aeruginosa*, *Staphylococcus epidermidis* (ATCC 35984), and *S. epidermidis* (ATCC 14990). In addition, Aumeeruddy *et al.* (2019) reported that its phenolic compounds could eliminate cancer cells (MCF-7). Table 2 shows the microorganism inhibited by the different types of honey from several countries.

The processing and storage conditions can impact antibacterial properties of honeys from different geographical and botanical origin against various pathogens, because some of them lose their bactericidal effect under 40-60 °C temperatures (Chen *et al.*, 2012; Pimentel González *et al.*, 2017; Matzen *et al.*, 2018).

Overview of the antibacterial properties of Mexican honey

Honey plays a very important role worldwide not just as food, but also as a medical-grade product. The five main exporters of honey are China, Argentina, Mexico, Germany, and Brazil (Campos *et al.*, 2018). Beekeeping has a major socioeconomic and ecologic importance in Mexico, because this farming activity generates many jobs (Magaña *et al.*, 2016). Honey is very appreciated in several countries, as a result of its nutritional properties, its perfume, taste, and color; consequently, Mexican honey is currently competing with honey produced in China, Vietnam, Nicaragua, Argentina, Chile, Turkey, and Ukraine (Martínez-González and Pérez-López, 2013; Campos *et al.*, 2018).

In Mexico, researches have included different topics: botanical characterization (Córdova-Córdova *et al.*, 2013), physicochemical properties, volatile compound profiles (Viuda-Martos *et al.*, 2010; Rodriguez *et al.*, 2012), color (Figure 2) (Grajales-Conesa *et al.*, 2018), and antioxidant activity of harvested honey (Leyva-Daniel *et al.*, 2017; Mondragón, 2019). However, few studies have focused on the antibacterial properties of honey and the identification of the main compounds that provide Mexican honey with the said properties (Maddock *et al.*, 2013).

Ramón-Sierra *et al.* (2020) compared the main phenolic compounds of honey produced by *Melipona beecheii* and *A. mellifera* from Merida, Yucatan. They found that the honey produced by *A. mellifera* had a higher phenolic compound and that 145 µg mL⁻¹ (phenolic extracts) and 60 µg mL⁻¹ (protein extracts) concentrations recorded antibacterial activities against *E. coli* and *S. aureus*. These results corroborated the findings of Chan-Rodríguez *et al.* (2012).

In addition, other studies have compared Mexican honey with honey from different countries. For example, some studies compare the honey of several honey bees (*Apis*) and the stingless bees from different countries such as Mexico (Jalisco, Chiapas, Quintana Roo, and Yucatan), Paraguay (Asuncion), Australia (Gatton and Brisbane), Philippines (Pili), Thailand (Chantaburi and Bangkok), Japan (Aichi), and Nepal. The evaluations were carried out to determine the effectiveness of those honeys against *Lactococcus lactis* ssp., *L. lactis cremoris* ssp. MAFF 40007, *Lactobacillus casei* JCM 1134, *E. faecalis* IFO 12964, *Enterococcus faecium* IFO 13712, *K. pneumoniae* ssp., and *Staphylococcus* ssp. The honeys from Yucatan and Chiapas (Mexico) recorded promising results in the inhibition of almost all the tested microorganisms, except for *K. pneumoniae* ssp. (Kimoto-Nira and Amano, 2008). Peláez-Acero *et al.* (2021) reported several new techniques to improve honey properties

Table 2. Minimum inhibitory concentration (MIC) of honeys from different geographical origins.

Country	floral source	Microorganisms that it inhibits	MIC (%)	Reference
Saudi Arabia	Dharm (<i>Lavandula dentata</i> L.), Majra (<i>Hypoestes forskaolii</i> (Vahl) R.Br.), Sider (<i>Ziziphus spina-christi</i> (L.) Willd)	<i>S. aureus</i> , <i>S. mutans</i> , <i>K. pneumoniae</i> , <i>E. coli</i> and <i>P. aeruginosa</i> .	20-30	Hegazi <i>et al.</i> , 2017.
Saudi Arabia	Yemeni Sidr and Mountain	Gram-negative bacteria	40-80	Alqurashi <i>et al.</i> , 2013
Algeria	Multifloral	<i>E. coli</i>	50-70	Bourabah <i>et al.</i> , 2020
Argentina	<i>Rapistrum rugosum</i> (L.) All., <i>Taraxacum officinale</i> F.H. Wigg.	<i>S. flexneri</i> , <i>Salmonella typhi</i> , <i>E. coli</i> .	100	Tejerina <i>et al.</i> , 2020
Australia	Jarrah (<i>Eucalyptus marginata</i> Donn ex Sm.) y Marri (<i>Corymbia calophylla</i> (Lindl.) KD Hill & LAS Johnson)	<i>Candida</i> spp. and dermatophytic fungi.	4-32	Irish <i>et al.</i> , 2011.
Chile	Ulmo (<i>Eucryphia cordifolia</i> Cav.)	<i>P. aeruginosa</i> , <i>S. aureus</i> , <i>E. coli</i> and <i>S. typhimurium</i>	9.4-20	Mullai y Menon, 2007; Sherlock <i>et al.</i> , 2010; Acevedo <i>et al.</i> , 2017; Velásquez <i>et al.</i> , 2020., Olate-Olave <i>et al.</i> , 2021
Denmark	Raspberry (<i>Rubus idaeus</i> L.), rapeseed (<i>Brassica napus</i> L.), Linden (<i>Tilia</i> L.) and Heather (<i>Calluna vulgaris</i> (L.) Hull)	<i>S. epidermidis</i> and <i>E. coli</i> .		Matzen <i>et al.</i> , 2018.
Ecuador	Multifloral	<i>S. aureus</i>	100	Montero-Recalde <i>et al.</i> , 2018
Egypt	Trefoil (<i>Trifolium repens</i> L.)	<i>S. typhimurium</i> and <i>E. coli</i> O157: H7	10-20	Mandal y Mandal, 2011.
Ethiopia	Multifloral	<i>S. aureus</i>	70-50	Mama <i>et al.</i> , 2019.
Ethiopia	Tazma mar (melipona), multiflorales	<i>S. aureus</i> , <i>P. aeruginosa</i> and <i>E. coli</i> .	9.38-11.25	Getaneh <i>et al.</i> , 2013.
Greece	Conifers, thyme (<i>Thymus</i> L.), citrus and multifloral	<i>S. typhimurium</i>	17-24	Voidarou <i>et al.</i> , 2011
India	Multifloral	<i>S. aureus</i> , <i>S. pyogenes</i> , <i>E. coli</i> , <i>P. aeruginosa</i> and <i>P. mirabilis</i>		Mahendran y Kumarasamy, 2015.
Mauritius Island	Eucalyptus (<i>Eucalipto globulus</i> Labill.)	<i>E. coli</i> ATCC 25922, <i>P. mirabilis</i> ATCC 12453, <i>P. aeruginosa</i> ATCC 27853, <i>S. epidermidis</i> ATCC35984, and <i>S. epidermidis</i> ATCC1499	100	Aumeeruddy <i>et al.</i> , 2019.
Malaysia	Tualang (<i>Koompassia excelsa</i> (Becc.) Taub.)	<i>E. coli</i> , <i>S. typhi</i> and <i>S. pyogenes</i> .	8.75-25	Mandal y Mandal, 2011.
Mexico	Acaxochitl (<i>Lobelia laxiflora</i> Kunth), multifloral	<i>B. subtilis</i> , <i>S. aureus</i> , <i>L. monocytogenes</i> , <i>E. coli</i> , <i>S. typhimurium</i> and <i>P. aeruginosa</i> .	55-100	Rodriguez <i>et al.</i> , 2012; Pimentel González <i>et al.</i> , 2017.
New Zealand	Manuka (<i>Leptospermum scoparium</i> J.R. Forst. and G. Forst.)	<i>S. pyogenes</i> , <i>S. mutans</i> , <i>P. mirabilis</i> , <i>P. aeruginosa</i> , <i>E. cloacae</i> and <i>S. aureus</i> .		Irish <i>et al.</i> , 2011; Johnston <i>et al.</i> , 2018.
Pakistan	Multifloral	<i>E. coli</i> , <i>P. aeruginosa</i> , <i>S. aureus</i> , <i>E. faecalis</i> .	75-100	Mustafa <i>et al.</i> , 2022
Portugal	Multifloral, <i>Leptospermum scoparium</i>	<i>E. coli</i> CECT 434, <i>E. coli</i> EC3a	25	Oliveira <i>et al.</i> , 2017
United Kingdom	Heather (<i>Calluna vulgaris</i>)	<i>S. aureus</i> , <i>E. coli</i> , <i>E. faecalis</i> and <i>P. vulgaris</i> .	2.5-40	Vučić <i>et al.</i> , 2014.



Figure 2. Color range of honey samples from Tabasco, Mexico.

(crystallization). These authors used an ultrasound to study the effect of liquefaction in the phenolic compounds, flavonoids, and antibacterial activity of crystallized honey from three multifloral honeys from Hidalgo. Their antibacterial activities were tested against *S. typhimurium*, *B. subtilis*, *P. aeruginosa*, *L. monocytogenes*, *S. aureus* and *E. coli*. Honeydew honey recorded a higher gallic acid content and some multifloral samples increased their quercetin content; meanwhile, all the samples inhibited the tested microorganisms. In particular, the antibacterial activity against *S. typhimurium* increased by 13%.

Overall, monofloral honeys have a higher inhibition range against pathogens than multifloral honeys (Oelschlaegel *et al.*, 2012; Rodriguez *et al.*, 2012; Escriche *et al.*, 2014; Pimentel-González *et al.*, 2017; Aumeeruddy *et al.*, 2019; Olate-Olave *et al.*, 2021).

Limitations

Currently, the use of honey for therapeutic effects has increased. It can be used for wound healing and to treat gastrointestinal diseases and eye infections. Nevertheless, evaluating the quality of the available scientific evidence is fundamental. On the one hand, information about the sample size is limited; on the other hand, the methods used to identify the properties of honey and the sample extraction before chromatography are not standardized. In addition, processing methods (pasteurized or unpasteurized) should be taken into account and the range of pathogenic bacteria evaluated should be expanded.

CONCLUSIONS

The antibacterial activity of honey has been used since ancient times to treat different health problems. Studying honey has never been an easy task: it is a complex system with inherent characteristics and many chemical compounds that change depending on the types of honey and its botanical and geographical origin. Overall, honeys produced from specific botanic sources have a high antibacterial activity, as a consequence of the medicinal properties of the flower source that are transferred to the honey through the nectar.

The efforts of the scientists to determine the underlying mechanisms of the bactericidal effect of honey have paid off. Scientists have proved that the said activity is related to the pH level, the osmotic effect caused by the high sugar content, the hydrogen peroxide content, the methylglyoxal content, the phenolic compounds, and the flavonoids that can work together or individually, depending on the type of honey (floral source, geographical

origin, etc.). All these characteristics help to stop the growth of certain pathogens and, consequently, honey has been used in the health sector to treat several diseases. Nevertheless, the antibacterial activity must be confirmed to prove the medical grade of a given honey. Subsequently, optimal production, handling, harvesting, and storage procedures should be established, in order to guarantee the safety of the product —*i.e.*, honey should be free of pathogens and toxic compounds. Finally, the effects of the infection and disease treatment should be proved through clinical trials.

The protocols used to study the phenolic compounds are different and are related to the detector that will be used with the HPLC. This selection is directly related to the analytes that will be identified and the budget established for the research. New technologies have led to promising results, reducing costs and analysis time; however, new technologies should be focused on the analysis of the metabolites of honey.

Consequently, further research about the different components of honey should be carried out to find new alternatives that reduce or stop antibiotic resistance. These studies should result in the use of honey as a substitute or its combination with classic antibiotics. Nevertheless, establishing standardized methodologies among the scientific community will help to determine the antibacterial activity of honey, facilitating its analysis and expanding the research of this property among honeys of different botanical origins.

REFERENCES

- Martínez-González, E. G., & Pérez-López, I. (2013). La producción de miel en el trópico húmedo de México: avances y retos en la gestión de la innovación. UACH. CIESTAAM. 102 p.
- Hegazi, A., El-Moez, S. I. A., Abdou, A. M., & Allah, F. A. (2014). Antibacterial Activity of Some Types of Monofloral Honey Against *Clostridium acetobutylicum* and *Clostridium perfringens*. *Int. J. Curr. Microbiol. Appl. Sci.*, 3(9), 552-565.
- Bueno-Costa, F. M., Zambiasi, R. C., Bohmer, B. W., Chaves, F. C., da Silva, W. P., Zanusso, J. T., & Dutra, I. (2016). Antibacterial and antioxidant activity of honeys from the state of Rio Grande do Sul, Brazil. *LWT - Food Sci. Tech-nol.*, 65(2016), 333-340. <https://doi.org/10.1016/j.lwt.2015.08.018>.
- Visweswara, R. P., Thevan, K. K., Salleh, N., & Hua G. S. (2016). Biological and therapeutic effects of honey produced by honey bees and stingless bees: a comparative review. *Rev. bras. farmacogn.*, 26(5), 657-664. <https://doi.org/10.1016/j.bjfp.2016.01.012>.
- Deng, J., Liu, R., Lu, Q., Hao, P., Xu, A., Zhang, J., & Tan, J. (2018). Biochemical properties, antibacterial and cellular antioxidant activities of buckwheat honey in comparison to manuka honey. *Food Chem.*, 252(2018), 243-249. <https://doi.org/10.1016/j.foodchem.2018.01.115>.
- Cheung, Y., Meenu, M., Yu, X., & Xu, B. (2019). Phenolic acids and flavonoids profiles of commercial honey from different floral sources and geographic sources. *Int. J. Food Prop.*, 22(1), 290-308. <https://doi.org/10.1080/10942912.2019.1579835>.
- Leyva-Jiménez, F. J., Lozano-Sánchez, J., Borras-Linares, I., Cadiz-Gurrea, M. L., & Mahmoodi-Khaledi, E. (2019). Potential antimicrobial activity of honey phenolic compounds against Gram positive and Gram-negative bacteria. *LWT - Food Sci. Technol.*, 101(2019), 236-245. <https://doi.org/10.1016/j.lwt.2018.11.015>.
- Alvarez-Suarez, J. M., Gasparrini, M., Forbes-Hernández, T. Y., Mazzoni, L., & Giampieri, F. (2014). The composition and biological activity of honey: a focus on Manuka honey. *Foods*, 3(3), 420-432. <https://doi.org/10.3390/foods3030420>.
- Güneş, M. E., Şahin, S., Demir, C., Borum, E., & Tosunoğlu, A. (2016). Determination of phenolic compounds profile in chestnut and floral honeys and their antioxidant and antimicrobial activities. *J. Food Biochem.*, 41(3), 1-12. <https://doi.org/10.1111/jfbc.12345>.
- Mandal, M. D., & Mandal, S. (2011). Honey: its medicinal property and antibacterial activity. *Asian Pac. J. Trop. Biomed.*, 1(2), 154-160. [https://doi.org/10.1016/S22211691\(11\)60016-6](https://doi.org/10.1016/S22211691(11)60016-6).
- Da Silva, G. J., & Mendonça, N. (2012). Association between antimicrobial resistance and virulence in *Escherichia coli*. *Virulence*, 3(1), 18-28. <https://doi.org/10.4161/viru.3.1.18382>.

- Fyfe, L., Okoro, P., Paterson, E., Coyle, S., & McDougall, G. J. (2017). Compositional analysis of Scottish honeys with antimicrobial activity against antibiotic-resistant bacteria reveals novel antimicrobial components. *LWT - Food Sci. Technol.*, 79(2017), 52-59. <https://doi.org/10.1016/j.lwt.2017.01.023>.
- Estrada, H., del Mar Gamboa, M., Chaves, C., & Arias, M. L. (2005). Evaluación de la actividad antimicrobiana de la miel de abeja contra *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Pseudomonas aeruginosa*, *Escherichia coli*, *Salmo-nella enteritidis*, *Listeria monocytogenes* y *Aspergillus niger*. Evaluación de su carga microbiológica. *Arch. Latinoam. Nutr.*, 55(2), 167-171.
- Boussaid, A., Chouaibi, M., Rezig, L., Hellal, R., Donsì, F., Ferrari, G., & Hamdi, S. (2018). Physicochemical and bio-active properties of six honey samples from various floral origins from Tunisia. *Arab. J. Chem.*, 11(2), 265-274. <https://doi.org/10.1016/j.arabjc.2014.08.011>.
- Jenkins, R. E., & Cooper, R. (2012). Synergy between oxacillin and manuka honey sensitizes methicillin-resistant *Staphylococcus aureus* to oxacillin. *J. Antimicrob. Chemother.*, 67(6), 1405-1407. <https://doi.org/10.1093/jac/dks071>.
- Müller, P., Alber, D. G., Turnbull, L., Schlothauer, R. C., Carter, D. A., Whitchurch, C. B., & Harry, E. J. (2013). Synergism between Medihoney and rifampicin against methicillin-resistant *Staphylococcus aureus* (MRSA). *PloS one*, 8(2), 1-9. <https://doi.org/10.1371/journal.pone.0057679>.
- Fernandes, L., Oliveira, A., Henriques, M., & Rodrigues, M. E. (2020). Honey as a Strategy to Fight *Candida tropicalis* in Mixed-Biofilms with *Pseudomonas aeruginosa*. *Antibiotics*, 9(2), 43-55. <https://doi.org/10.3390/antibiotics9020043>.
- Goslinski, M., Nowak, D., & Klebukowska, L. (2020). Antioxidant properties and antimicrobial activity of manuka honey versus Polish honeys. *J. Food Sci. Technol.* 57(4), 1269-1277. <https://doi.org/10.1007/s13197-019-04159-w>.
- Velásquez, P., Montenegro, G., Leyton, F., Ascar, L., Ramirez, O., & Giordano, A. (2020). Bioactive compounds and antibacterial properties of monofloral Ulmo honey. *CYTA - J. Food.*, 18(1), 11-19. <https://doi.org/10.1080/19476337.2019.1701559>.
- Almasaudi, S. B., Al-Nahari, A. A., El Sayed, M., Barbour, E., Al Muayawi, S. M., Al-Jaoumi, S., Mohamad, Q., Yousef, A. Q., & Harakeh, S. (2017). Antimicrobial effect of different types of honey on *Staphylococcus aureus*. *Saudi J. Biol. Sci.*, 24(6), 1255-1261. <https://doi.org/10.1016/j.sjbs.2016.08.007>.
- Wen-Jie, N. G., & Mei-Siew, L. (2015). Anti-staphylococcal activity of Melaleuca honey. *Southeast Asian J. Trop. Med. Public Health*, 46(3), 472-479.
- Schencke, C., Vásquez, B., Sandoval, C., & del Sol, M. (2016). El rol de la miel en los procesos morfofisiológicos de reparación de heridas. *Int. J. Morphol.*, 34(1), 385-395. <http://dx.doi.org/10.4067/S0717-95022016000100056>.
- Irish, J., Blair, S., & Carter, D. A. (2011). The antibacterial activity of honey derived from Australian flora. *Plos one*, 6(3), e18229. <https://doi.org/10.1371/journal.pone.0018229>.
- Habib, H. M., Al Meqbali, F. T., Kamal, H., Souka, U. D., & Ibrahim, W. H. (2014). Physicochemical and biochemical properties of honeys from arid regions. *Food Chem.*, 153(2014): 35-43. <https://doi.org/10.1016/j.foodchem.2013.12.048>.
- Johnston, M., McBride, M., Dahiya, D., Owusu-Apenten, R., & Nigam, P. S. (2018). Antibacterial activity of Manuka honey and its components: An overview. *AIMS Microbiol.*, 4(4), 655-662. <https://doi.org/10.3934%2Fmicrobiol.2018.4.655>.
- Matzen, R. D., Zinck Leth-Espensen, J., Jansson, T., Nielsen, D. S., Lund, M. N., & Matzen, S. (2018). The antibacterial effect *in vitro* of honey derived from various Danish flora. *Dermatol. Res. Pract.*, 021713(2018), <https://doi.org/10.1155/2018/7021713>.
- Atrott, J., & Henle, T. (2009). Methylglyoxal in manuka honey-correlation with antibacterial properties. *Czech J. Food Sci.*, 27(2009), S163-S165. <https://doi.org/10.17221/911-CJFS>.
- Adams, C. J., Manley-Harris, M., & Molan, P. C. (2009). The origin of methylglyoxal in New Zealand manuka (*Leptospermum scoparium*) honey. *Carbohydr. Res.*, 344(8), 1050-1053. <https://doi.org/10.1016/j.carres.2009.03.020>.
- Kwakman, P. H., & Zaat, S. A. (2012). Antibacterial components of honey. *IUBMB life*, 64(1), 48-55. <https://doi.org/10.1002/iub.578>.
- Pharma GDD. 2022. Revamil: du miel médical. Recuperado el 15 de abril de 2023. Disponible en: <https://www.pharma-gdd.com/fr/revamil>.
- Iqbal, M. N., Anjum, A. A., Ali, M. A., Hussain, F., Ali, S., Muhammad, A., Irfan, M., Ahmad, A., Irfan, M., & Shabbir, A. (2015). Assessment of microbial load of unpasteurized fruit juices and *in vitro* antibacterial potential of honey against bacterial isolates. *Open Microbiol. J.*, 9(2015), 26-32. <https://doi.org/10.2174%2F1874285801509010026>.

- Albaridi, N. A. (2019). Antibacterial potency of honey. *Int. J. Microbiol.*, 2464507(2019), 1-10. <https://doi.org/10.1155/2019/2464507>.
- Mama, M., Teshome, T., & Detamo, J. (2019). Antibacterial activity of honey against methicillin-resistant *Staphylococcus aureus*: a laboratory-based experimental study. *Int. J. Microbiol.*, 7686130(2019) 1-9. <https://doi.org/10.1155/2019/7686130>.
- Cembrero, G., Sanhueza, O., Pezoa, M., Báez, M. E., Martínez, J., Báez, M., & Fuentes, E. (2020). Relationship among the minor constituents, antibacterial activity and geographical origin of honey: A multifactor perspective. *Food Chem.*, 315(2019), 1-9. <https://doi.org/10.1016/j.foodchem.2020.126296>.
- Cauich, K. R., Ruiz, R. J. C., Ortíz, V. E., & Segura, C. M. R. (2015). Potencial antioxidante de la miel de *Melipona beecheii* y su relación con la salud: una revisión. *Nutr. Hosp.*, 32(4), 1432-1442. <https://dx.doi.org/10.3305/nh.2015.32.4.9312>.
- Hossen, M. S., Ali, M. Y., Jahurul, M. H. A., Abdel-Daim, M. M., Gan, S. H., & Khalil, M. I. (2017). Beneficiales roles of honey polyphenols against some human degenerative diseases: A review. *Pharmacol. Rep.*, 69(6): 1194-1205. <https://doi.org/10.1016/j.pharep.2017.07.002>.
- Brudzynski, K., & Miotto, D. (2011). The relationship between the content of Maillard reaction-like products and bioactivity of Canadian honeys. *Food Chem.*, 124(3): 869-874. <https://doi.org/10.1016/j.foodchem.2010.07.009>.
- Sant'Ana, L. D. O., Buarque Ferreira, A. B., Lorenzon, M. C. A., Berbara, R. L. L., & Castro, R. N. (2014). Correlation of total phenolic and flavonoid contents of Brazilian honeys with colour and antioxidant capacity. *Int. J. Food Prop.*, 17(1), 65-76. <https://doi.org/10.1080/10942912.2011.614368>.
- Sousa, J. M., De Souza, E. L., Marques, G., Meireles, B., de Magalhães Cordeiro, Â. T., Gullón, B., Pintado, M. M. & Magnani, M. (2016). Polyphenolic profile and antioxidant and antibacterial activities of monofloral honeys produced by Meliponini in the Brazilian semiarid region. *Food Res. Int.*, 84(2016), 61-68. <https://doi.org/10.1016/j.foodres.2016.03.012>.
- Escuredo, O., Silva, L. R., Valentão, P., Seijo, M. C., & Andrade, P. B. (2012). Assessing Rubus honey value: Pollen and phenolic compounds content and antibacterial capacity. *Food Chem.*, 130(3), 671-678. <https://doi.org/10.1016/j.foodchem.2011.07.107>.
- Poli, J. P., Guinoiseau, E., Luciani, A., Yang, Y., Battesti, M. J., Paolini, J., Costa, Y., Quilichini, L., & Lorenzi, V. (2018). Key role of hydrogen peroxide in antimicrobial activity of spring, Honeydew maquis and chestnut grove Corsican honeys on *Pseudomonas aeruginosa* DNA. *Lett. Appl. Microbiol.*, 66(5), 427-433. <https://doi.org/10.1111/lam.12868>.
- Yi, H. Y., Chowdhury, M., Huang, Y. D., & Yu, X. Q. (2014). Insect antimicrobial peptides and their applications. *Appl. Microbiol. Biotechnol.*, 98(13), 5807-5822. <https://doi.org/10.1007/s00253-014-5792-6>.
- Majtan, J., Klaudiny, J., Bohova, J., Kohutova, L., Dzurova, M., Sediva, M., Bartosova, M., & Majtan, V. (2012). Methyl-glyox-al-induced modifications of significant honeybee proteinous components in manuka honey: possible therapeutic implications. *Fitoterapia*, 83(4), 671-677. <https://doi.org/10.1016/j.fitote.2012.02.002>.
- Dosler, S., & Karaaslan, E. (2014). Inhibition and destruction of *Pseudomonas aeruginosa* biofilms by antibiotics and antimicrobial peptides. *Peptides*, 62(2014): 32-37. <https://doi.org/10.1016/j.peptides.2014.09.021>.
- Valachová, I., Bučeková, M., & Majtán, J. (2016). Quantification of Bee-Derived Peptide Defensin-1 in Honey by Competitive Enzyme-Linked Immunosorbent Assay, a New Approach in Honey Quality Control. *Czech J. Food Sci.*, 34(3), 233-243. <http://dx.doi.org/10.17221/422/2015-CJFS>.
- Čerovský, V., & Bém, R. (2014). Lucifensins, the insect defensins of biomedical importance: the story behind maggot therapy. *Pharmaceuticals*, 7(3), 251-264. <https://doi.org/10.3390/ph7030251>.
- Larsen, A., Reynaldi, F. J., & Guzmán-Novoa, E. (2019). Bases del sistema inmune de la abeja melífera (*Apis mellifera*). Revisión. *Rev. Mex. Cienc. Pecu.*, 10(3), 705-728. <https://doi.org/10.22319/rmcv.v10i3.4785>.
- Li, Y., Xiang, Q., Zhang, Q., Huang, Y., & Su, Z. (2012). Overview on the recent study of antimicrobial peptides: origins, functions, relative mechanisms and application. *Peptides*, 37(2), 207-215. <https://doi.org/10.1016/j.peptides.2012.07.001>.
- Danihlík, J., Aronstein, K., & Petřivalský, M. (2015). Antimicrobial peptides: a key component of honey bee innate immunity: Physiology, biochemistry, and chemical ecology. *J. Apic. Res.*, 54(2), 123-136. <https://doi.org/10.1080/00218839.2015.1109919>.
- Randolt, K., Gimple, O., Geissendörfer, J., Reinders, J., Prusko, C., Mueller, M. J., & Beier, H. (2008). Immune related proteins induced in the hemolymph after aseptic and septic injury differ in honey bee worker larvae and adults. *Arch. Insect Biochem. Physiol.*, 69(4): 155-167. <https://doi.org/10.1002/arch.20269>.
- Erban, T., Shcherbachenko, E., Talacko, P., & Harant, K. (2019). The unique protein composition of honey revealed by comprehensive proteomic analysis: allergens, venom-like proteins, antibacterial properties,

- royal jelly proteins, serine proteases, and their inhibitors. *J. Nat. Prod.*, 82(5), 1217-1226. <https://doi.org/10.1021/acs.jnatprod.8b00968>.
- Khalil, A. T., Khan, I., Ahmad, K., Khan, Y. A., Khan, J., & Shinwari, Z. K. (2014). Antibacterial activity of honey in north-west Pakistan against select human pathogens. *J. Tradit. Chin. Med.*, 34(1), 86-89. [https://doi.org/10.1016/S0254-6272\(14\)60059-5](https://doi.org/10.1016/S0254-6272(14)60059-5).
- Osés, S. M., Pascual-Mate, A., de la Fuente, D., de Pablo, A., Fernandez-Muino, M. A., & Sancho, M. T. (2016). Comparison of methods to determine antibacterial activity of honeys against *Staphylococcus aureus*. *NJAS - Wageningen J. Life Sci.*, 78(2016), 29-33. <https://doi.org/10.1016/j.njas.2015.12.005>.
- Balouiri, M., Sadiki, M., & Ibnsouda, S. K. (2016). Methods for *in vitro* evaluating antimicrobial activity: A review. *J. Pharm. Anal.*, 6(2), 71-79. <https://doi.org/10.1016/j.jpha.2015.11.005>.
- Clavijo, D. A. (2002). Fundamentos de química analítica: equilibrio iónico y análisis químico. Univ. Nacional de Colombia. Bogotá, Colombia. 977pp.
- Camaró-Sala, M. L., Martínez-García, R., Olmos-Martínez, P., Catalá-Cuenca, V., Ocete-Mochón, M. D., & Gime-no-Cardona, C. (2015). Validación y verificación analítica de los métodos microbiológicos. *Enferm. Infect. Microbiol. Clin.*, 33(7), 31-36. <https://doi.org/10.1016/j.eimc.2013.11.010>.
- Roshan, N., Rippers, T., Locher, C., & Hammer, K. A. (2017). Antibacterial activity and chemical characteristics of several Western Australian honeys compared to Manuka honey and pasture honey. *Arch. Microbiol.*, 199(2), 347-355. <https://doi.org/10.1007/s00203-016-1308-3>.
- Rosas, C. D. O., Rodrigues, J. M., Cruz, M. H. C. D. L., Lopes, S. M. D. R., Souto, A. D. S. S., Brandão, M. L. L., & Ca-passo, I. R. V. D. F. (2019). Microbiological reference material (bacterial and fungal domains): definition, production rules, use and need for establishment in Brazil. *Braz. J. Food Technol.*, 22(2019), 1-11. <https://doi.org/10.1590/1981-6723.20817>.
- Pasiás, I. N., Kiriakou, I. K., Kaitatzis, A., Koutelidakis, A. E., & Proestos, C. (2018). Effect of late harvest and floral origin on honey antibacterial properties and quality parameters. *Food Chem.*, 242(2018), 513-518. <https://doi.org/10.1016/j.foodchem.2017.09.083>.
- Hegazi, A. G., Al Guthami, F. M., Al Gethami, A. F., Abd Allah, F. M., Saleh, A. A., & Fouad, E. A. (2017). Potential antibacterial activity of some Saudi Arabia honey. *Vet. World*, 10(2), 233-237. <https://doi.org/10.14202%2Fvetworld.2017.233-237>.
- Mahendran, S., & Kumarasamy, D. (2015). Antimicrobial activity of some honey samples against pathogenic bacteria. *Int. Lett. Nat. Sci.*, 7(2015), 15-20.
- Aumeeruddy, M. Z., Aumeeruddy-Elalfi, Z., Neetoo, H., Zengin, G., van Staden, A. B., Fibrich, B., Lambrechts, I. A., Rademan, S., Szuman, K. M., Lall, N., & Mahomoodally, F. (2019). Pharmacological activities, chemical profile, and physicochemical properties of raw and commercial honey. *Biocatal. Agric. Biotechnol.*, 18(2019), 1-10. <https://doi.org/10.1016/j.biocab.2019.01.043>.
- Chen, C., Campbell, T.L., Blair, E.S., & Carter, A.D. (2012). The effect of standard heat and filtration processing procedures on antimicrobial activity and hydrogen peroxide levels in honey. *Front. Microbiol.*, 3(2012), 1-8. <https://doi.org/10.3389/fmicb.2012.00265>.
- Alqurashi, A.M., Masoud, E.A., & Alamin, M.A., 2013. Antibacterial activity of Saudi honey against Gram negative bacteria. *J. Microbiol. Antimicrob.*, 5, 1-5. <https://doi.org/10.5897/JMA2012.0235>.
- Bourabah, A., Titri, A., Heni, K., Ahmad, M., & Aissa, A., 2020. Evaluation of antibacterial activity and additive effect between casein and two types of algerian honey. *Adv. Biol. Earth Sci.*, 5, 128-134.
- Tejerina, M. R., Cabana, M. J., Sánchez, A. C., & Benítez, A. M. R., 2020. Antibacterial effect of argentinian pollens and honeys. *J. Microbiol. Biotech. Food Sci.* 10, 381-384. <https://doi.org/10.15414/jmbfs.2020.10.3.381-384>.
- Mullai, V., & Menon, T. (2007). Bactericidal activity of different types of honey against clinical and environmental isolates of *Pseudomonas aeruginosa*. *J. Altern. Complement. Med.*, 13(4), 439-442. <https://doi.org/10.1089/acm.2007.6366>.
- Sherlock, O., Dolan, A., Athman, R., Power, A., Gethin, G., Cowman, S., & Humphreys, H., 2010. Comparison of the antimicrobial activity of Ulmo honey from Chile and Manuka honey against methicillin-resistant *Staphylococcus aureus*, *Escherichia coli* and *Pseudomonas aeruginosa*. *BMC Complement. Altern. Med.*, 10, 1-5. <https://doi.org/10.1186/1472-6882-10-47>.
- Acevedo, F., Torres, P., Oomah, B. D., de Alencar, S. M., Massarioli, A. P., Martín-Venegas, R., Albarral-Ávila, V., Ferrer, R., & Rubilar, M., 2017. Volatile and non-volatile/semi-volatile compounds and *in vitro* bioactive properties of Chilean Ulmo (*Eucryphia cordifolia* Cav.) honey. *Food Res. Int.*, 94, 20-28. <https://doi.org/10.1016/j.foodres.2017.01.021>.
- Olate-Olave, V. R., Guzmán, L., López-Cortés, X. A., Cornejo, R., Nachtigall, F. M., Doorn, M., Silva, S. L., & Bejarano, A., 2021. Comparison of Chilean honeys through MALDI-TOF-MS profiling and

- evaluation of their antioxidant and antibacterial potential. *Ann. Agric. Sci.*, 66, 152-161. <https://doi.org/10.1016/j.aoas.2021.11.001>.
- Montero-Recalde, M., Vayas, L., Avilés-Esquivel, D., Pazmiño, P., & Erazo-Gutierrez, V., 2018. Evaluación de dos métodos para medir la sensibilidad de inhibición de crecimiento de la cepa certificada de *Staphylococcus aureus* subsp. *aureus*. *Rev. de Investig. Vet. del Perú*, 29, 1543-1547.
- Getaneh, A., Belyhun, Y., Moges, F., Anagaw, B., Destaw, B., Unakal, C., & Mulu, A. (2013). *In vitro* assessment of the antimicrobial effect of Ethiopian multiflora honey on methicillin resistant *Staphylococcus aureus*. *Int. J. Cur. Res. Rev.*, 5(11), 64-72.
- Voidarou, C., Alexopoulos, A., Plessas, S., Karapanou, A., Mantzourani, I., Stavropoulou, E., Fotou, F., Tzora, A., Skoufos, I., & Bezirtzoglou, E., 2011. Antibacterial activity of different honeys against pathogenic bacteria. *Anaerobe*, 17, 375-379. <https://doi.org/10.1016/j.anaerobe.2011.03.012>.
- Rodríguez, B. A., Mendoza, S., Iturriga, M. H., & Castaño Tostado, E. (2012). Quality parameters and antioxidant and antibacterial properties of some Mexican honeys. *J. Food Sci.*, 77(1), C121-C127. <https://doi.org/10.1111/j.1750-3841.2011.02487.x>.
- Pimentel González, D.J., Basilio Cortes, U. A., Hernández Fuentes, A. D., Figueira, A. C., Quintero Lira, A., & Cam-pos Montiel, R. G. (2017). Effect of thermal processing on antibacterial activity of multifloral honeys. *J. Food Process Eng.*, 40(1), e12279. <https://doi.org/10.1111/jfpe.12279>.
- Mustafa, G., Iqbal, A., Javid, A., Manzoor, M., Aslam, S., Ali, A., Azam, S. M., Khalid, M., Farooq, M., Al-Naggar, Y., Ali-Alharbi, S., Ali-El, E. H., Abd-Malek, R., Qamer, S., & Hussain, A., 2022. Antibacterial properties of *Apis dorsata* honey against some bacterial pathogens. *Saudi J. Biol. Sci.*, 29, 730-734. <https://doi.org/10.1016/j.sjbs.2021.09.059>.
- Oliveira, A., Ribeiro, H. G., Silva, A. C., Silva, M. D., Sousa, J. C., Rodrigues, C. F., Melo, L. D. R., Henriques, A. F., & Sillankorva, S., 2017. Synergistic antimicrobial interaction between honey and phage against *Escherichia coli* biofilms. *Front. Microbiol.*, 8, 1-18. <https://doi.org/10.3389/fmicb.2017.02407>.
- Vučić, D. M., Petković, M. R., Rodić-Grabovac, B. B., Stefanović, O. D., Vasić, S. M., & Čomić, L. R. (2014). *In vitro* activity of heather [*Calluna vulgaris* (L.) Hull] extracts on selected urinary tract pathogens. *Bosn. J. Basic Med. Sci.*, 14(4), 234-238. <https://doi.org/10.17305%2Fbjbms.2014.4.40>.
- Pyrzynska, K., & Biesaga, M. (2009). Analysis of phenolic acids and flavonoids in honey. *TrAC, Trends Anal. Chem.*, 28(7), 893-902. <https://doi.org/10.1016/j.trac.2009.03.015>.
- Ciulu, M., Spano, N., Pilo, M. I., & Sanna, G. (2016). Recent advances in the analysis of phenolic compounds in unifloral honeys. *Molecules*, 21(4), 1-32. <https://doi.org/10.3390/molecules21040451>.
- Sowa, P., Grabek-Lejko, D., Wesołowska, M., Swacha, S., & Dzugan, M. (2017). Hydrogen peroxide-dependent antibacterial action of *Melilotus albus* honey. *Lett. Appl. Microbiol.* 65(1), 82-89. <https://doi.org/10.1111/lam.12749>.
- Sindi, A., Chawn, M.V.B., Hernández, E.M., Green, K., Islam, K.M., Locher, C., & Hammer, K. (2019). Anti-biofilm effects and characterisation of the hydrogen peroxide activity of a range of Western Australian honeys compared to Manuka and multifloral honeys. *Sci. Rep.* 9(2019), 1-17. <https://doi.org/10.1038/s41598-019-54217-8>.
- Sultambawa, Y., Cozzolino, D., Fuller, S., Cusack, A., Currie, M., & Smyth, H. (2015). Infrared spectroscopy as a rapid tool to detect methylglyoxal and antibacterial activity in Australian honeys. *Food Chem.* 172(2015), 207-212. <https://doi.org/10.1016/j.foodchem.2014.09.067>.
- Cokcetin, N.N., Pappalardo, M., Campbell, L.T., Brooks, P., Carter, D.A., Blair, S.E., & Harry, E.J. (2016). The Antibacterial Activity of Australian Leptospermum Honey Correlates with Methylglyoxal Levels. *PLOS ONE*, 11(12), 1-13. <https://doi.org/10.1371/journal.pone.0167780>.
- Rückriemen, J., Klemm, O., & Henle, T. (2017). Manuka honey (*Leptospermum scoparium*) inhibits jack bean urease activity due to methylglyoxal and dihydroxyacetone. *Food Chem.*, 230(2017), 540-546. <https://doi.org/10.1016/j.foodchem.2017.03.075>.
- Escríche, I., Kadar, M., Juan-Borrás, M., & Domenech, E. (2014). Suitability of antioxidant capacity, flavonoids and phenolic acids for floral authentication of honey. Impact of industrial thermal treatment. *Food Chem.*, 142(2014), 135-143. <https://doi.org/10.1016/j.foodchem.2013.07.033>.
- Ouchemoukh, S., Amessis-Ouchemoukh, N., Gómez-Romero, M., Aboud, F., Giuseppe, A., Fernández-Gutiérrez, A., & Segura-Carretero, A. (2017). Characterisation of phenolic compounds in Algerian honeys by RP-HPLC coupled to electrospray time-of-flight mass spectrometry. *LWT - Food Sci. Technol.*, 85(2017), 460-469. <https://doi.org/10.1016/j.lwt.2016.11.084>.
- Elrasheid, T. H., Xiaoobo, Z., Zhihua, L., Jiyoung, S., Zhai, X., Wang, S., & Mariod, A. A. (2017). Rapid prediction of phenolic compounds and antioxidant activity of Sudanese honey using Raman and Fourier transform infrared (FT-IR) spectroscopy. *Food Chem.*, 226(2017), 202-211. <https://doi.org/10.1016/j.foodchem.2017.01.024>.

- Stanek, N., & Jasicka-Misiak, I. (2018). HPTLC phenolic profiles as useful tools for the authentication of honey. *Food Anal. Methods*, 11(2018), 2979-2989. <https://doi.org/10.1007/s12161-018-1281-3>.
- Gośliński, M., Nowak, D., & Szwengiel, A. (2021). Multidimensional comparative analysis of bioactive phenolic compounds of honeys of various origin. *Antioxidants*, 10(4), 530. <https://doi.org/10.3390/antiox10040530>.
- Stanek, N., Kafarski, P., & Jasicka-Misiak, I. (2019). Development of a high performances thin layer chromatography method for the rapid qualification and quantification of phenolic compounds and abscisic acid in honeys. *J. Chro-matogr. A.* 1598(2019), 209-215. <https://doi.org/10.1016/j.chroma.2019.04.052>.
- Moniruzzaman, M., Yung An, C., Rao, P. V., Hawlader, M. N. I., Azlan, S. A. B. M., Sulaiman, S. A., & Gan, S. H. (2014). Identification of Phenolic Acids and Flavonoids in Monofloral Honey from Bangladesh by High Performance Liquid Chromatography: Determination of Antioxidant Capacity. *Biomed. Res. Int.*, 37490(2014), 1-11. <https://doi.org/10.1155/2014/737490>.
- Montenegro, G., Gómez, M., Diaz-Forestier, J., & Pizarro, R. (2008). Aplicación de la Norma Chilena Oficial de denominación de origen botánico de la miel para la caracterización de la producción apícola. *Cienc. Inv. Agr.*, 35(1): 181-190. <http://dx.doi.org/10.4067/S0718-16202008000200007>.
- Silva, L. R., Gonçalves, A. C., Nunes, A. R., & Alves, G. (2020). Authentication of honeys from Caramulo region (Portugal): Pollen spectrum, physicochemical characteristics, mineral content, and phenolic profile. *Journal of food science*, 85(2), 374-385. <https://doi.org/10.1111/1750-3841.15023>.
- Badjah H. A., A. Y., Wabaidur, S. M., Siddiqui, M. R., Alothman, Z. A., Obeid, M. S., Khan, M. R., & Altamrah, S. A. (2016). Simultaneous determination of twenty-five polyphenols in multifloral and cactus honeys using solid-phase extraction and high-performance liquid chromatography with photodiode array detection. *Eur. Food Res. Technol.*, 242(6), 943-952. <https://doi.org/10.1007/s00217-015-2601-0>.
- Tang, K. S., Konczak, I., & Zhao, J. (2016). Identification and quantification of phenolics in Australian native mint (*Mentha australis* R. Br.). *Food Chem.*, 192(2016), 698-705. <https://doi.org/10.1016/j.foodchem.2015.07.032>.
- Mani, J. S., Johnson, J. B., Hosking, H., Ashwath, N., Walsh, K. B., Neilsen, P. M., Broszczak, D., & Naiker, M. (2021). Antioxidative and therapeutic potential of selected Australian plants: A review. *Journal of Ethnopharmacology*, 268(2021), 1-16. <https://doi.org/10.1016/j.jep.2020.113580>.
- Campos, G. M., Leyva, M. C., Ferráez, P. M., & Sánchez, B. Y. (2018). El mercado internacional de la miel de abeja y la competitividad de México. *Rev. econ.*, 35(90), 87-123.
- Magaña, M. M. A., Tavera, C. M. E., Salazar, B. L. L., & Sanginés, G. J. R. (2016). Productividad de la apicultura en México y su impacto sobre la rentabilidad. *Rev. Mex. Cienc. Agríc.*, 7(5), 1103-1115.
- Córdova-Córdova, C. I., Ramírez-Arriaga, E., Martínez-Hernández, E., y Zaldívar-Cruz, J. M. (2013). Caracterización botánica de miel de abeja (*Apis mellifera* L.) de cuatro regiones del estado de Tabasco, México, mediante técnicas melisopalinológicas. *Universidad y ciencia*, 29(2): 163-178.
- Viuda Martos, M., Ruiz Navajas, Y., Zaldívar Cruz, J. M., Kuri, V., Fernández López, J., Carbonell Barrachina, Á. A., & Pérez Álvarez, J. (2010). Aroma profile and physicochemical properties of artisanal honey from Tabasco, Mexico. *Int. J. Food Sci. Technol.*, 45(6), 1111-1118. <https://doi.org/10.1111/j.1365-2621.2010.02243.x>.
- Leyva-Daniel, D. E., Escobedo-Avellaneda, Z., Villalobos-Castillejos, F., Alamilla-Beltrán, L., & Welti-Chanes, J. (2017). Effect of high hydrostatic pressure applied to a Mexican honey to increase its microbiological and functional quality. *Food Bioprod. Process.*, 102(2017), 299-306. <https://doi.org/10.1016/j.fbp.2017.01.001>.
- Mondragón, C. P. (2019). Antioxidant Capacity and Total Phenolic Content in Honey Brands from Mexican Market and Some Physicochemical Parameters Related. *World J. Food Sci. Technol.*, 3(2), 20-25. <https://doi.org/10.11648/j.wjfst.20190302.11>.
- Maddock, S. E., Jenkins, R. E., Rowlands, R. S., Purdy, K. J., & Cooper, R. A. (2013). Manuka honey inhibits adhesion and invasion of medically important wound bacteria *in vitro*. *Future Microbiol.*, 8(12), 1523-1536. <https://doi.org/10.2217/fmb.13.126>.
- Grajales-Conesa, J., Vandame, R., Santiesteban-Hernández, A., López-García, A., & Guzmán-Díaz, M. (2018). Propiedades fisicoquímicas y antibacterianas de mieles de abejas sin aguijón del sur de Chiapas, México. *Ibciencias*, 1(1), 1-7.
- Ramón-Sierra, J., Martínez-Guevara, J. L., Pool-Yam, L., Magana-Ortiz, D., Yam-Puc, A., & Ortiz-Vázquez, E. (2020). Effects of phenolic and protein extracts from *Melipona beecheii* honey on pathogenic strains of *Escherichia coli* and *Staphylococcus aureus*. *Food Sci. Biotechnol.*, 29(7), 1013-1021. <https://doi.org/10.1007/s10068-020-00744-4>.

- Chan-Rodríguez, D., Ramón-Sierra, J., Lope-Ayora, J., Sauri-Duch, E., Cuevas-Glory, L., & Ortiz-Vázquez, E. (2012). Antibacterial properties of honey produced by *Melipona beecheii* and *Apis mellifera* against foodborn microorganisms. *Food Sci. Biotechnol.*, 21(3), 905-909. <https://doi.org/10.1007/s10068-012-0118-x>.
- Kimoto-Nira, H., & Amano, K. (2008). Antimicrobial activity of honey produced by stingless honey bees. *J. Apic. Res.*, 47(4), 325-327. <https://doi.org/10.1080/00218839.2008.11101484>.
- Peláez-Acero, A., Cobos-Velasco, J. E., González-Lemus, U., Espino-Manzano, S. O., Aguirre-Álvarez, G., González-Montiel, L., & Campos-Montiel, R. G. (2021). Bioactive compounds and antibacterial activities in crystallized honey liquefied with ultrasound. *Ultrason Sonochem.*, 76(2021), 1-6. <https://doi.org/10.1016/j.ulsonch.2021.105619>.
- Oelschlaegel, S., Gruner, M., Wang, P. N., Boettcher, A., Koelling-Speer, I., & Speer, K. (2012). Classification and Characterization of Manuka Honeys Based on Phenolic Compounds and Methylglyoxal. *J. Agric. Food Chem.*, 60(29), 7229-7237. <https://doi.org/10.1021/jf300888q>.

